

An Introduction to Sedges


Richard Carter

Herbarium

Biology Department

Valdosta State University

Valdosta, GA 31698



Cyperaceae – the sedge family

- Third largest monocot family
- ~5000 species, 104 genera
- Largest genera
 - *Carex*, 2000 spp.
 - *Cyperus*, 550 spp. (excl. *Kyllinga*, *Pycneus*)
 - *Fimbristylis*, 300 spp.
 - *Rhynchospora* and *Scleria*, 250 spp. each
 - *Eleocharis*, 200 spp.
 - *Bulbostylis*, *Pycneus* and *Schoenus*, 100 spp. each

General features of sedges


- Grass-like, monocot flowering plants
- Linear leaves, parallel venation
- Small, mostly wind-pollinated flowers



This is not a sedge, but it is a monocot!

Note perianth of 6 parts.

Lilium catesbaei Walter
pine lily



Phylogenetic relationships


- Traditionally viewed as close relatives of the grasses (Poaceae)
- Recent cladistic analysis using molecular & morphological data shows closer alliance with Juncaceae & Thurniaceae.

(3,4)

Graminoids



- Sedges, grasses, rushes and other similar kinds of monocot plants with small, inconspicuous flowers and linear leaves are grouped informally as graminoids.






Sedges have edges....

- *Sedges have edges; rushes are round; grasses are hollow right up from the ground.*
- Most sedges have 3-angled stems, hence *sedges have edges*; however, some do not.
 - E.g., stems of *Dulichium arundinaceum* and many *Eleocharis* species are round (terete) in cross section.




Comparison of grasses, rushes & sedges

<i>Cyperaceae</i> The Sedge Family	<i>Poaceae</i> The Grass Family	<i>Juncaceae</i> The Rush Family
• Stems usually three-angled (but sometimes terete, quadrangular, or lenticular)	• Stems terete	• Stems terete
• Stems usually with solid pith	• Stems with solid nodes and hollow internodes	• Stems with solid pith
• Leaf sheaths closed	• Leaf sheaths open	• Leaf sheaths open
• Inflorescence a complex of spikelets (simple spikelet in <i>Eleocharis</i>)	• Inflorescence a complex of spikelets	• Inflorescence a complex of cymes
• Perianth of 1—many bristles or hairs, or absent	• Perianth hardly evident, apparently reduced to scale-like palea (outer series?) and tiny lodicule (inner series)	• Perianth of six scale-like parts in two series
• Stamens 3 (1-2, rarely 6)	• Stamens 3 or 6 (rarely 1-2)	• Stamens 6 (rarely 3)
• Pistil of 2-3 fused carpels	• Pistil of 2(3) fused carpels	• Pistil of 3 fused carpels
• Fruit an achene	• Fruit a caryopsis (grain)	• Fruit a capsule



Common names can be confusing!

- Many graminoids, sedges included, escape all but passing notice and do not have common names.
- Common names are often derived uncritically.
 - Bulrushes (*Scirpus* spp., *Schoenoplectus* spp.), spike-rushes (*Eleocharis* spp.), and beak-rushes (*Rhynchospora* spp.) are sedges.
 - Cotton-grasses (*Eriophorum* spp.), umbrella-grasses (*Fuirena* spp.), and sawgrass (*Cladium jamaicense*) are sedges.
 - The nut-sedges (*Cyperus esculentus*, *C. rotundus*) are often called “nut-grasses.”
- Being universal & unambiguous, scientific names promote precise communication.




Ancient uses of sedges

- Papyrus (*Cyperus papyrus*)
 - First exploited by ancient Egyptians ~4500 years ago to manufacture paper
 - English word “paper” from Latin name for this species
- A bulrush, *Schoenoplectus corymbosus*, used in funeral wreaths

Cyperus papyrus L. cultivated in water garden




San Diego County,
California, USA



Food for humans

- Chufas
 - Tubers from *Cyperus esculentus* var. *sativus*, yellow-nutsedge
 - One of the oldest crops in Egypt
 - Cultivated in Africa, Asia and southern Europe
 - Rich in starch, sugar and fat
 - Nutty flavor when roasted
 - Can be made into flour
 - Spanish drink *horchata de chufas*
 - Source of non-drying oil of some economic value
- Chinese water-chestnut
 - Tubers of aquatic spikerush, *Eleocharis dulcis*
 - Grown in paddies in Asia
- Rhizomes of bulrushes (*Schoenoplectus* spp.) were eaten by native Americans


(7,8)



Food for wildlife

- Tubers of yellow nut-sedge and other sedges are eaten by wildlife
- Fruits (achenes) of aquatic sedges *Eleocharis*, *Schoenoplectus*, etc. consumed by waterfowl

(9,10)




Sedges as ornamentals

- Umbrella sedge (*Cyperus alternifolius* subsp. *flabelliformis*) has been grown in water gardens and as a pot-plant for more than 200 years!
- Water gardens & ponds
 - papyrus (*Cyperus papyrus*)
 - dwarf papyrus (*Cyperus prolifer* Kunth)
 - bulrushes (*Scirpus* spp., *Schoenoplectus* spp.)
- Woodland gardens
 - *Carex* spp.
- Potted plants & hanging baskets
 - *Cyperus albostriatus*
 - *Isolepis cernuus*

(11,12)



Cyperus involucratus Rottb.
cultivated in water garden
Lowndes County, Georgia, USA




Miscellaneous uses

- Robust bulrushes, like *Schoenoplectus californicus* (C.A. Mey.) Soják, exploited to construct houses and boats
- Stems, leaves, or fibers of many sedges used as materials for weaving, especially in undeveloped parts of the world
 - E.g., stems and leaves of various bulrushes (*Scirpus* spp., *Schoenoplectus* spp.) are woven into baskets, mats, and chair seats
 - *Scirpus americanus* Pers. commonly called chairmaker's rush
 - Fibers from *Fimbristylis umbellaris* (Lam.) Vahl used as material for weaving in Asia
- Water purification
 - Bulrush *Schoenoplectus lacustris* (L.) Palla in Germany & the Netherlands
- Indicators of copper deposits
 - *Fimbristylis* spp. in Australia

(2,7)

Sedges have long been recognized among world's worst agricultural weeds.



1st *Cyperus rotundus* L.

16th *Cyperus esculentus* L.

32nd *Cyperus difformis* L.

33rd *Cyperus iria* L.

40th *Fimbristylis miliacea* (L.) Vahl

The world's worst weed!


Cyperus rotundus L. purple nutsedge

Aggressive perennial weed of
agricultural & urban areas


- Prolific production of rhizomes & tubers
 - Seed rarely produced
 - Rapid growth
 - Allelopathy
 - C₄ photosynthesis
- (13)



Purple nutsedge in cotton




Purple nutsedge growth in 60 days – Dr. Wills




Sedges can be taxonomically challenging!

- Extreme reduction of flowers and fruits in size and number
- Inherent difficulty in handling and describing such small, specialized parts
 - Good hand lens or dissecting microscope required
 - Ability to manipulate and dissect fine structures
- Reliable identification requires reproductively mature specimens with fully developed spikelets and achenes.



General Structure

- Habit
- Leaves and stems
- Inflorescence
- Flowers
- Fruits and associated structures




Habit

- Annual or perennial herbs
- Mostly perennial herbs persisting and spreading vegetatively by rhizomes, stolons, corms, or tubers

Diminutive annual

Cyperus pumilus L.

Clinch Co., Georgia



Cespitose perennial

Eleocharis tuberculosa (Michaux) R. & S.
Margin of drawn down flatwoods pond,
Atkinson Co., Georgia, USA





Vegetative proliferation by rhizomes & tubers

Eleocharis acutangula (Roxb.) Schult.

Lee County, Florida




“Walking” vegetative proliferation of aerial stems

Eleocharis melanocarpa Torr.

Nyssa biflora-*Taxodium ascendens*-*Ilex myrtifolia*-*Litsea* pond

Turner Co., Georgia



Aerial stems and leaves

- Stems typically trigonous – 3 sides, 3 angles
 - Exceptions previously noted
- Leaves
 - Arise at intervals along leafy stem (e.g., *Dulichium*, *Scirpus*, *Bolboschoenus*)
 - Clustered near base of plant (e.g., *Cyperus*, *Kyllinga*)
 - Closed sheathing bases
 - Blades lanceolate to linear, grass-like
 - Parallel venation
- Unique within the family, plants of *Eleocharis* have leaves reduced to bladeless sheaths and, thus, appear leafless.




Closed leaf sheath
Lanceolate blade

Dulichium arundinaceum (L.) Britt.



Parallel venation

Cymophyllus fraserianus (Ker-
Gawl.) Kartesz & Gandhi



Inflorescences

- Some genera, e.g., *Cyperus* and *Kyllinga*, have prominent leafy bracts subtending inflorescence.
- Spikelet = basic unit of inflorescence
- Organization of spikelets in inflorescence varies
 - Paniculate
 - Cymose
 - Umbellate
 - Spicate
 - Solitary

A close-up photograph of a plant stem, likely a grass or sedge, showing a spiral arrangement of floral scales. The stem is green and has a series of brown, overlapping scales that spiral upwards. The background is a blurred field of similar plants.

Spiral arrangement of floral scales

Eleocharis equisetoides (Ell.) Torr.

Clinch Co., Georgia

Distichous arrangement of floral scales
Cyperus sanguinolentus Vahl



1.0 mm



Umbellate inflorescence
subtended by leafy bracts

Cyperus strigosus L.

Baker Co., Florida

Spikes and spikelets



Cyperus croceus Vahl




Intact spikelet
and spikelet with portion of floral
scale removed to show fruit (inset)

br



spk




pr




Cyperus croceus Vahl

General structure



Cyperus retrofractus (L.) Torr.



Flowers

- Wind-pollinated (anemophilous)
- Highly reduced both in size and numbers of parts
- Usually perfect
 - Imperfect – *Carex*, *Cymophyllus*, *Scleria*
- Androecium
 - 1-3 stamens
- Gynoecium
 - 2-3 carpellate pistil
- Perianth
 - Extremely reduced or completely absent
 - When present, perianth usually persists attached to fruit
 - Bristles – animal dispersal (zoochory)
 - e.g., *Eleocharis*, *Rhynchospora*
 - Hairs – wind dispersal (anemochory)
 - e.g., *Scirpus*, *Eriophorum*
 - Bristles + paddle-shaped segments
 - *Fuirena*

Flowering spikelets



Eleocharis montevidensis Kunth

Grady County, Georgia



Gynoecium – pistil

- 3-carpellate, 3-branched (trifid) style
- 2-carpellate, 2-branched (bifid) style




Anemophily

Fimbristylis puberula
(Michx.) Vahl

Flowers generally
protogynous

Exposed feathery,
stigmas promote
wind pollination






Stigmas projecting
beyond perigynium
Carex striata Michx.



Complex perianth


Fuirena breviseta (Cov.) Cov.



1.5kV

X60 200µm 0000 29 30 SEI

Achenes and associated structures




- Achene = sedge fruit
 - Small
 - 1-seeded
 - Dry
 - Indehiscent
- Mature achenes essential for reliable identification of species
 - Shape, size, color, surface ornamentation taxonomically useful
- Achene shape correlated with carpel number
 - Pistils derived from 3 carpels have 3-branched (trifid) styles and form trigonous or terete achenes
 - Pistils with 2 carpels normally have 2-branched (bifid) styles and develop into biconvex (lenticular) or plano-convex achenes.

Achene shape

correlated with number of carpels
& style branches

- Trigonous: 3-carpellate pistil, 3-branched style
- Biconvex (lenticular): 2-carpellate pistil, 2-branched style





Achene-tubercle- perianth complex


Eleocharis tuberculosa
(Michx.) R. & S.

Toothed perianth bristles
promote dispersal of achenes by
attachment to hair & feathers

Function of spongy tubercle

Buoyancy? – water dispersal?

Lipid? – dispersal by ants?




Wind dispersal by persistent, silky perianth

Scirpus cyperinus (L.) Kunth




Survey of the major groups of sedges



- Spike-rush Sedge Group
- Bulrush Sedge Group
- Umbrella-grass Sedge Group
- Fringe-sedge Group
- Flat-sedge Group
- Three-Way Sedge Group
- Beak-rush Sedge Group
- Sawgrass Sedge Group
- Nut-rush Sedge Group
- Caric Sedge Group


Generic classification follows FNA (10).

Etymology of generic names
(10,15,16)



The Spike-rush Sedge Group

- Diagnostic characteristics
 - Plants apparently leafless, with bladeless leaves reduced to sheathing bases
 - Inflorescence a single, terminal, unbranched spikelet
 - Flowers perfect
- 2 genera
 - *Eleocharis*
 - *Websteria*



Eleocharis – Spike-rushes

- From Greek *elos*, marsh, and *charis*, grace
- Most structurally reduced sedges, consisting of little more than an apparently leafless stem terminated by simple spikelet
- Taxonomy based largely on characteristics of perianth, tubercles, surface ornamentation of achenes
- Mostly on hydric soils
 - Wetlands
 - Floodplains
 - Seasonally wet sites in fields and pastures


Eleocharis tuberculosa
(Michx.) R. & S.
Baker Co., Florida



Endozoic dispersal of achenes by waterfowl

Eleocharis equisetoides (Ell.) Torr.

Clinch Co., Georgia, USA



Mature spikelet, just
before separation of
achenes



Eleocharis acutangula (Roxb.) Schult.
Florida, USA




Vegetative proliferation
Eleocharis baldwinii (Torr.) Chapm.
Lowndes Co., Georgia, USA





Eleocharis montevidensis Kunth
Grady Co., Georgia



Websteria

- Commemorating G. W. Webster, American botanist and farmer, 1833-1914
- Monotypic genus – *Websteria confervoides* (Poir.) Hooper
- Diagnostic characteristics
 - Vegetatively similar to *Eleocharis vivipara* Link.
 - Submerged aquatic
 - Stems capillary
 - Stems forming false whorls
 - Scales distichous
 - Spikelets 1-fruited
- Habitat and distribution
 - Widely distributed in tropical, subtropical and warm temperate regions around the world
 - United States
 - Infrequently collected
 - Known only from Florida and Georgia
 - Submersed in ponds and lakes

(10,17)

Websteria confervooides (Poir.) Hooper


Lake Co., Florida

Submerged plants (background),
dislodged floating plants (inset)




Photographs courtesy of
Nia Wellendorf, Florida DEP

The Bulrush Sedge Group



- Diagnostic characteristics
 - Scales spiral
 - Flowers perfect
 - Perianth of bristles or hairs, or absent
 - Style base indistinct
 - Tubercle absent
- Segregate genera traditionally included in *Scirpus* or Scirpeae
 - *Bolboschoenus*
 - *Schoenoplectus*
 - *Isolepis**
 - *Oxycaryum**
 - *Lipocarpha**

*More recently allied with Cyperae (1,10,18)



Scirpus – Bulrushes

- Classical Latin name for the bulrush
- Diagnostic characteristics
 - Leafy stems
 - Large, compound, cymose inflorescence of many spikelets
 - Scales
 - Glabrous
 - Usually acute to acuminate tips
- Various habitats, on hydric soils

A close-up photograph of the seed heads of Scirpus cyperinus. The image shows several dense, elongated panicles of small, brownish, fuzzy seed heads hanging from thin green stems. The background is a soft-focus green, suggesting a natural outdoor setting.

Scirpus cyperinus (L.) Kunth
Woolly bulrush, woolly bully
McIntosh Co., Georgia




Dispersal along roads & railroads
Scirpus cyperinus (L.) Kunth
Wayne Co., Georgia

Scirpus divaricatus Ell.

Camden Co., Georgia





Schoenoplectus – Naked-stem Bulrushes


- From Greek, *schoinos*, rush, and *plectos*, plaited, referring to use of stems in weaving of mats, etc.
- Diagnostic characteristics
 - Leafless, wand-like stems
 - Ciliate scales
- ~10 species in Georgia, e.g.
 - *Schoenoplectus etuberculatus* (Steud.) Soják
 - Emergent in shallow ponds of the coastal plain or laxly submersed in swiftly flowing blackwater streams
 - *Schoenoplectus pungens* (Vahl) Palla
 - Coastal salt-marsh species
 - With pseudolateral clusters of sessile spikelets subtended by an erect bract that appears to be a continuation of stem



Schoenoplectus etuberculatus (Steud.) Soják
Berrien Co., Georgia


Schoenoplectus etuberculatus (Steud.) Soják
Berrien Co., Georgia





Schoenoplectus pungens (Vahl) Palla
McIntosh Co., Georgia





Bolboschoenus – Tuberous Bulrushes


- Greek *bolbos*, bulb, and *schoinos*, rush, referring to enlarged, cormous stem bases
- Diagnostic characteristics
 - Cormous stem bases
 - Leafy stems
 - Large spikelets
 - Puberulent scales

Bolboschoenus robustus (Pursh) Soják
Cameron Co., Texas



Bolboschoenus robustus (Pursh) Soják
Cameron Co., Texas





Isolepis


- From Greek, *isos*, equal, and *lepis*, scale, referring to the uniform floral scales
- Diagnostic characteristics
 - Low, cespitose habit
 - Basal leaves
 - Terminal or pseudolateral capitate or solitary inflorescences
- Recent molecular evidence indicates closer relationship with Cyperaceae


(1,4,10,18)

Isolepis

- 2 spp. SE United States, both annuals
 - *I. carinata* Hook. ex Arn. ex Torr. [= *Scirpus koilolepis* Steud.]
 - Native
 - Spring ephemeral of intermittently wet depressions of fields and open woods
 - *I. pseudosetacea* (Dav.) Gand. [= *Scirpus molestus* M.C. Johnst.]
 - Introduction
 - Similar habitat and phenology

Source of illustration: Ball, W, K Gandhi, RW Kiger, D Murray, JL Zarucchi, AA Reznicek and JL Strother. 2002. *Flora of North America*, vol. 23, Oxford University Press. New York.





Oxycaryum

- From Greek, *oxys*, sharp, and *carya*, nut, referring to the sharp-pointed achene
- Monotypic genus – *Oxycaryum cubense* (Poepp. & Kunth) Lye [= *Scirpus cubensis* Poepp. & Kunth]
- Recent molecular evidence indicates closer relationship with Cyperaceae
- Diagnostic characteristics
 - Stoloniferous, floating aquatic
 - Terminal, umbellate or monocephalous inflorescence
 - Subtended by whorl of leafy bracts
 - Resembles *Cyperus* or *Kyllinga*
 - Spiral scales

(1,4,10,17,18)




Oxycaryum cubense
fo. *paraguayense* (Maury) Pedersen

monocephalous form

Achene with corky pericarp,
dispersed by water




Oxycaryum cubense (Poepp. & Kunth) Palla



Impounded bayswamp
Georgia, USA




- Broad, paleotropical & neotropical distribution
- Perennial, spreading locally by stolons, forming extensive floating mats in swamps & ponds
- Known from SE USA pre-1900 – TX, LA, s AL, FL
- Currently spreading in SE USA
 - 1996 – s GA
 - 2004 – Tennessee-Tombigbee River system in MS & AL



Lipocarpha

- Classification of *Lipocarpha* depends on how one interprets the various kinds of scales in the inflorescence and, thus, whether one views the inflorescence as a simple spikelet or a compound spike.
 - Simple spikelet – Scirpeae
 - Compound spike – Cyperae
- Recent molecular evidence supports classification in Cyperae
- Conundrum illustrates struggle inherent in two fundamental purposes of taxonomy to provide stable and ultimately useful means of identifying and naming plants and to construct classification schemes that reflect phylogenetic (evolutionary) relationships

(1,4,10,18)



Lipocarpha

- From Greek *leipo*, to fall, and *carpha*, chaff, referring to the deciduous inner scales of certain species
- *Lipocarpha maculata* (Michx.) Torr.
 - Resembles *Kyllinga* with its cespitose habit and terminal inflorescence of tightly clustered spikelets subtended by a whorl of leafy bracts
 - Distribution and habitat
 - Occasional to common in the coastal plain
 - Wet ditches, disturbed hydric soils of depressions in the flatwoods, and along the exposed margins of ponds


Lipocarpa maculata (Michx.) Torr.
Charlton Co., Georgia



The Umbrella-grass Sedge Group

- Diagnostic characteristics
 - Leaf blades or sheaths usually pubescent
 - Scales spiral, usually pubescent
 - Flowers perfect
 - Perianth differentiated into two series, 3 outer bristles and 3 inner paddle-like segments
 - Achene with stipitate base and peg-like apex
 - Tubercle absent






Fuirena – Umbrella-grasses

- Commemorating Georg Fuiren, Danish Botanist, 1581-1628
- 5 spp. in SE United States
 - *F. breviseta* (Cov.) Cov.
 - *F. longa* Chapm.
 - *F. pumila* (Torr.) Spreng.
 - *F. scirpoidea* Michx.
 - *F. squarrosa* Michx.
- Habitat – heliophytes of wetland habitats, including bogs, marshes, interdunal swales, ditches, margins of ponds, and wet depressions in savannas

Fuirena breviseta (Cov.) Cov.

Clinch Co., Georgia





The Fringe-sedge Group

- Diagnostic characteristics
 - Leaves basal
 - Inflorescences terminal
 - Scales spiral
 - Flowers perfect
 - Perianth absent
 - Style-base distinct
 - Tubercle present (*Bulbostylis*) or absent (*Fimbristylis*)
- 3 genera
 - *Abildgaardia*
 - *Fimbristylis*
 - *Bulbostylis*

Fimbristylis – Fringe-sedges




- From Latin *fimbria*, fringed, and *stylus*, style
- Diagnostic characteristics
 - Tubercle absent
 - Fringed style with base clearly distinct from summit of achene
 - Branched, umbellate inflorescence of several to many spikelets
- Some widely distributed weeds introduced from Old World via rice agriculture
 - *F. annua* (All.) R. & S.
 - *F. dichotoma* (L.) Vahl
 - *F. miliacea* (L.) Vahl
 - *F. tomentosa* Vahl
- Species of conservation concern
 - *F. perpusilla* Harper
 - *F. brevivaginata* Kral

(10,20,21,22,23)

Introduced with rice agriculture

Fimbristylis miliacea (L.) Vahl



- Widely distributed in tropical & warm temperate regions of E & W Hemispheres
- Common weed of rice
- Probably indigenous to Asian rice belt
- Numerous small seed (10,13,20)





Fimbristylis autumnalis (L.) R. & S.
Bacon Co., Georgia

Bulbostylis


(10,20)



- From Latin *bulbus*, bulbous, and *stylus*, style, referring to the enlarged bulbous style bases of many species
- Diagnostic characteristics
 - Swollen style base forming distinct tubercle on summit of achene
- Habitat
 - Seasonally moist to xeric sands
- *Bulbostylis barbata* (Rottb.) C. B. Clarke
 - Diminutive annual
 - Reddish-brown inflorescences
 - Open, disturbed sandy loam
 - Conspicuous en masse in the coastal plain during late summer and autumn
 - Widespread in E and W Hemispheres
- *Bulbostylis warei* (Torrey) C. B. Clarke
 - Cespitose perennial
 - Hemispherical, head-like clusters of spikelets
 - Distinctive inflorescence bracts with beautifully fringed basal sheaths
 - Endemic to Atlantic and Gulf coastal plains of SE United States
 - Inhabits open sands in longleaf pine-scrub oak communities


Bulbostylis barbata (Rottb.) Clarke





The Flat-sedge Group

- Diagnostic characteristics
 - Leaves basal
 - Leafy bracts subtending inflorescence
 - Inflorescence terminal, umbellate with pedunculate rays or capitate cluster of sessile spikes
 - Scales distichous
 - Flowers perfect
 - Perianth absent
 - Style base indistinct
 - Tubercle absent
- Largest, most taxonomically complex group



Cyperus –

Flat-sedges or Umbrella Sedges

- From Greek *cyperus*, edge, referring to the sharp-edged leaves or perhaps the three-edged stems
- Diagnostic characteristics
 - Floral scales distichous (2-ranked)
 - Usually 2+ flowers or fruits per spikelet
 - Perianth absent
- Large complex genus – 550-700 spp.

Cyperus

Classification of subgenera

- Basis of classification
 - Leaf anatomy and photosynthetic pathway
 - Number of carpels & style branches
 - Achene shape
 - Achene orientation
 - Mode of spikelet disarticulation & unit of dispersal
- Subgenera
 - Stigmas 3, achenes trigonous
 - *Anosporum* (Nees) Clarke [= *Pycnostachys* Clarke]
 - *Cyperus* L.
 - *Diclidium* (Schrad. Ex Nees) Clarke [= *Torulium* (Desv. ex Ham.) Kük.]
 - Stigmas 2, achenes biconvex
 - *Pycneus* (Beauv.) Gray
 - *Juncellus* (Griseb.) Clarke

C₃ & C₄ photosynthesis


C₄ photosynthesis, kranz anatomy

- Many agricultural weeds
- Lower CO₂ compensation point
- Increased water use efficiency
- Plants more competitive
 - at higher ambient temperatures
 - during drought




C₃ photosynthesis, non-kranz anatomy

- Fewer weeds
- Plants generally adapted to hydric or mesic environments
- May be competitive in other ways in hydric or mesic environments



C₃ & C₄ photosynthesis among weedy sedge genera

(12,18,26,27)



Achene shape

correlated with style branch number

- Trigonous: *Cyperus*,
Anosporum, *Diclidium*
- Lenticular (biconvex):
Kyllinga, *Pycneus*,
Juncellus



Achene orientation lenticular achenes only


- Angle adjacent to rachilla: *Pycneus*
- Face adjacent to rachilla: *Juncellus*



C. laevigatus

Dispersal of individual achenes

Floral scales and achenes separate sequentially from base to apex of spikelet rachilla.



Cyperus haspan L.

Dispersal of
entire spikelet



Cyperus echinatus (L.) Wood



--Spikelet breaking transversely into 1-2 fruited segments


--Water dispersal by corky rachilla



Cyperus odoratus L.


Cyperus cuspidatus Kunth
Lanier Co., Georgia






Cyperus echinatus (L.) Wood
Lowndes Co., Georgia





Cyperus retrorsus Chapm.
Lowndes Co., Georgia





Cyperus nashii Britt. ex Small

Marion Co., Florida



Epizoaic dispersal of spikelet
with pungent terminal scale
Cyperus plukenetii Fern.



Subgenus *Diclidium*

Cyperus odoratus L.


McIntosh Co., Georgia



Subgenus *Anosporum*


Cyperus difformis L.

Copious production of small
achenes, short generation time (13)



Subgenus *Pycreus*

Cyperus sanguinolentus Vahl




- Introduced from Asia – rice weed
- Currently, expanding its range in SEUS
- Habitat – disturbed sites, e.g., roadside ditches, margins of ponds
- Annual, small achenes
- Dispersed by highway maintenance equipment




Subgenus *Juncellus*

Cyperus alopecuroides Rottb.
recently naturalized in Florida



- Native – paleotropics
- Naturalized – West Indies & Florida, USA
- Perennial
- Numerous, small achenes
- Invasive tendencies, forming floating mats in reclamation wetlands
- Potential threat to natural wetlands & limesink ponds in Florida



Kyllinga – Green Sedges

- Name commemorating Peter Kylling, 17th century Danish botanist
- Diagnostic characteristics
 - Terminal, capitate inflorescence
 - 2-scaled, 1-fruited spikelets
 - Lenticular achenes
- *Kyllinga* closely related to and probably derived from *Cyperus*, and sometimes treated within *Cyperus* as a subgenus or section
- Five species in SE United States
 - *K. brevifolia* Rottb. – In US before 1821
 - *K. gracillima* Miq.
 - *K. odorata* Vahl – In US before 1836
 - *K. pumila* Michx. – In US before 1803, native?
 - *K. squamulata* Thonn. ex Vahl



Dispersal of entire spikelet
Kyllinga odorata Vahl




1 mm



Kyllinga odorata Vahl
Lowndes Co., Georgia




Kyllinga squamulata Thonn. ex Vahl
Lowndes Co., Georgia



The Three-Way Sedge Group

- Diagnostic characteristics
 - Stem terete
 - Leaves cauline
 - Upper leaves with well-developed lanceolate blades, conspicuously three-ranked
 - Inflorescences axillary
 - Scales distichous
 - Flowers perfect
 - Perianth of 6-9 bristles



Dulichium – Three-way Sedge


- Latin name for a kind of sedge
- Monotypic genus – *Dulichium arundinaceum* (L.) Britt.
- Combination of perianth bristles and distichous scales found elsewhere among sedges in SE United States only in *Eleocharis baldwinii* and *Websteria confervoides*
- Habitat – acidic soils of depressions along blackwater streams and shallows along ponds associated with such streams



Dulichium arundinaceum (L.) Britt.
Hamilton Co., Florida




The Beak-rush Sedge Group



- Diagnostic characteristics
 - Scales spiral
 - Flowers perfect
 - Perianth of few to many bristles or absent in sections *Dichromena* and *Psilocarya*
 - Stigmas 2 (-3)
 - Achene biconvex to subterete
 - Tubercle present

Rhynchospora – Beak-rushes



- From Greek *rhyncho*, snout or beak, and *spora*, seed, referring to the beaked achenes of many species
- Most beak-rushes inhabit hydric soils in bogs, wet savannas, margins of ponds, seeps, and depressions in flatwoods
 - *R. megalocarpa* Gray and *R. grayi* Kunth found in open, xeric, sandy pinelands or sandscrub
- Some, opportunistic colonizers of pastures, lawns, pond margins, and ditches, are treated as weeds
 - *R. caduca* Ell. recently naturalized, spreading rapidly in Hawaii
- Beak-rushes of conservation concern include
 - *R. crinipes* Gale – banks and bars of blackwater streams
 - *R. harveyi* var. *culixa* (Gale) Kral – ecotones between sandhills and bogs
 - *R. solitaria* Harper – hillside bogs
 - *R. thornei* Kral – margins of limesink ponds

(10,22,32,33)

Achene – tubercle – perianth

Rhynchospora inexpansa (Michx.) Vahl



Rhynchospora miliacea (Lam.) Gray
Cook Co., Georgia



Rhynchospora cephalantha Gray
Brooks Co., Georgia



Rhynchospora ciliaris (Michx.) Mohr
Charlton Co., Georgia



Rhynchospora macrostachya
Torr. ex Gray
Pierce Co., Georgia



Rhynchospora

Section *Dichromena*

Section *Psilocarya*

- Section *Dichromena* – conspicuous dichromatic white and green inflorescence bracts
 - *R. colorata* (L.) Pfeiff. – basic to circumneutral soils in seeps or swales
 - *R. latifolia* (Baldw.) Thomas – acidic soils of bogs and wet savannas
 - *R. floridensis* (Britt. ex Small) Pfeiff. – solution pits in limerock in S Florida
- Section *Psilocarya* – annuals lacking perianth bristles
 - *R. nitens* (Vahl) Gray
 - *R. scirpoides* (Torr.) Gray

Section *Dichromena*


Rhynchospora colorata (L.) Pfeiff.

Lanier Co., Georgia



Section *Psilocarya*
Rhynchospora nitens (Vahl) Gray
Baker Co., Florida





Schoenus – black sedge


- From Greek *schoinos*, rush-like plant
- Allied with *Rhynchospora*, usually placed in separate tribe Schoeneae
- Diagnostic characteristics
 - Inflorescence terminal to pseudolateral, capitate
 - Scales distichous, black
 - Perianth bristles usually 6, short, basally sub-plumose
 - Achene whitish
 - Tubercle absent
- Seasonally wet calcareous outcrops

Schoenus nigricans L. – black sedge
calcareous glade, Gadsden Co., Florida




Schoenus nigricans L. – black sedge
calcareous glade, Gadsden Co., Florida





The Sawgrass Sedge Group

- Diagnostic characteristics
 - Scales spiral
 - Flowers perfect
 - Perianth absent
 - Stigmas 3
 - Achene terete




Cladium – Sawgrass

- From Greek *clados*, branch, alluding to the branched inflorescence
- Only two species in SE United States
 - *C. jamaicense* Crantz
 - Inhabits brackish and freshwater marshes along the Georgia coast and occasionally inland
 - Dominant species of Everglades marshes of S Florida
 - Robust perennial
 - Graceful, delicate inflorescences
 - Lacerating foliage
 - *C. mariscoides* (Muhl.) Torrey


Cladium jamaicense Crantz
Franklin Co., Florida




Inflorescence of Sawgrass



The Nut-rush Sedge Group




- Diagnostic characteristics
 - Flowers imperfect
 - Spikelet generally with pistillate flowers below staminate and with several empty basal scales
 - Achenes whitish, bony
 - Hypogynium often present



Scleria – Nut-rushes

- From Greek *scleros*, hard, referring to the bony achene
- Features of achene and hypogynium taxonomically useful
 - Whitish, bony surfaces of achenes smooth, pitted, reticulate or pubescent
 - Hypogynium (usually present) fused to base of achene – discoid, tuberculate, or lobed
- Habitat
 - Most species on fairly wet sites, e.g., open, moist, sandy or peaty soils of seepage slopes, bogs, depressions in flatwoods, and pond margins
 - *S. triglomerata* Michx. and *S. oligantha* Michx. more often on mesic to subxeric sites in shaded woods, open prairies, and pineland savannas
 - *S. ciliata* Michx. and *S. pauciflora* Muhl. ex Willd. both exhibit ample variation with several named varieties each and substantial ranges in habitat from dry to hydric sites





Scleria reticularis Michx.
netted nut-rush







Three-lobed hypogynium and netted achene surface
in *Scleria reticularis* Michx.



Scleria reticularis Michx.



Aculeate hypogynium
Scleria triglomerata Michx.




1.5KV

X35 500µm 0000 28 30 SEI




Hypogynium elevated on broad inverted, cuplike base
Scleria oligantha Michx.






Hypogynium absent
Scleria georgiana Core

The Caric Sedge Group



- Diagnostic characteristics
 - Flowers imperfect
 - Plants usually monoecious (dioecious in *Carex picta*)
 - Staminate (male) and pistillate (female) flowers often borne in separate inflorescences or one type above the other in the same inflorescence
 - Sac-like perigynium enclosing each pistillate flower and achene
- 2 genera in SE United States
 - *Carex*
 - *Cymophyllus* – monotypic



Carex


- From Greek *cairo*, to cut, referring to sharp edges of leaves in certain species
- Habitat, etc.
 - Mostly in mesic, woodland habitats of northern temperate zone
 - >2000 species
 - Largest genus of Cyperaceae
 - One of the largest genera of the world's flora

Flowers imperfect

Staminate & pistillate spikes separate

Carex glaucescens Ell.





Flowers imperfect
Staminate flowers below
pistillate in same spike
Carex sect. *Ovales*

A close-up photograph of a pistillate spike of Carex striata. The spike is green and shows several perigynia, which are the female flower clusters. Each perigynium is surrounded by a bracteole, a protective leaf-like structure. The perigynia are arranged in a spiral pattern along the spike. The bracteoles are reddish-brown and have a distinct V-shaped notch at their tips. The spike is covered in fine, white, hair-like structures. The background is a blurred green leaf.

Portion of pistillate
spike showing
perigynia and
bracteoles
Carex striata Michx.

Perigynium

Pistillate spikelet with
perigynium face cut
away, exposing
gynoecium within

--*Carex striata* Michx.






Carex lonchocarpa Willd. ex Spreng.
Lowndes Co., Georgia



Carex striata Michx.
Cypress-gum pond
Echols Co., Georgia




Carex tenax Chapm.
Turner Co., Georgia




Acknowledgements

- Faculty Development Fund, Valdosta State University
- Dr. Russell Goddard, Biology Department, Valdosta State University, assisted with scanning electron microscopy.
- Scanning electron microscopy was supported by National Science Foundation award DBI-0420454




References

1. Goetghebeur, P. 1998. Cyperaceae, pp. 141-190 in: K. Kubitzki (ed.), *The families and genera of vascular plants IV*. Springer-Verlag. Berlin.
2. Simpson, D. A. and C. A. Inglis. 2001. Cyperaceae of economic, ethnobotanical, and horticultural importance: a checklist. *Kew Bulletin* 56: 257-360.
3. Cronquist, A. 1981. *An integrated system of classification of flowering plants*. Columbia University Press. New York.
4. Chase, M. W., D. E. Soltis, P. S. Soltis, P. J. Rudall, M. F. Fay, W. H. Hahn, S. Sullivan, J. Joseph, M. Molvray, P. J. Kores, T. J. Givnish, K. J. Sytsma and J. C. Pires. 2000. Higher-level systematics of the monocotyledons: an assessment of current knowledge and a new classification, pp. 3-16 in: Wilson, K. L. and D. A. Morrison (eds.), *Monocots: systematics and evolution*. CSIRO Publishing. Collingwood, Victoria.
5. Carter, R. 2005. An introduction to the sedges of Georgia. *Tipularia* 20:17-45.
6. Levitin, E. and K. McMahon. 2003. *Plants and society*. McGraw-Hill Companies, Inc. New York.
7. Mabberley, D. J. 1997. *The plant book*. Cambridge University Press. Cambridge.
8. Schery, R. W. 1972. *Plants for man*. Prentice-Hall, Inc. Englewood Cliffs, New Jersey.
9. DeFelice, M. S. 2002. Yellow nutsedge *Cyperus esculentus* L. – snack food of the gods. *Weed Technol.* 16: 901-907.
10. Ball, P. W., K. Gandhi, R. W. Kiger, D. Murray, J. L. Zarucchi, A. A. Reznicek and J. L. Strother. 2002. *Flora of North America*, vol. 23, Oxford University Press. New York.
11. Bailey, L. H. and E. Z. Bailey. 1976. *Hortus third*. Macmillan Publishing Company, Inc. New York.



References

12. Bryson, C. T. and R. Carter. In press. The significance of Cyperaceae as weeds. Monog. Syst. Bot. Missouri Bot. Garden.
13. Holm, L.G., D. Plucknett, J. V. Pancho, and J. P. Herberger. 1977. *The world's worst weeds: distribution and biology*. University Press of Hawaii. Honolulu.
14. Terry, P. J. 2001. The Cyperaceae – still the world's worst weeds? Pages 3-18, in C. R. Riches. *The World's Worst Weeds*. British Crop Protection Council. Farnham.
15. Fernald, M. L. 1950. *Gray's manual of botany*. 8th edition (reprint). Dioscorides Press. Portland, Oregon.
16. Hyam, R. and R. Pankhurst. 1995. *Plants and their names: a concise dictionary*. Oxford University Press Inc. New York.
17. Godfrey, R. K. and J. W. Wooten. 1979. *Aquatic and wetland plants of southeastern United States: monocotyledons*. University of Georgia Press. Athens.
18. Bruhl, J. 1995. Sedge genera of the world: relationships and a new classification of the Cyperaceae. *Australian Systematic Botany* 8: 125-305.
19. Kral, R. 1978. A synopsis of *Fuirena* (Cyperaceae) for the Americas north of South America. *Sida* 7: 309-354.
20. Kral, R. 1971. A treatment of *Abildgaardia*, *Bulbostylis*, and *Fimbristylis* (Cyperaceae) for North America. *Sida* 4: 58-277.
21. Patrick, T. S., J. R. Allison and G. A. Krakow. 1995. *Protected plants of Georgia*. Georgia Department of Natural Resources. Social Circle.
22. Kral, R. 1983. *A report on some rare, threatened, or endangered forest-related vascular plants of the South: vol. 1, Isoetaceae through Euphorbiaceae*. Technical Publication R8-TP 2. USDA Forest Service. Atlanta, Georgia.



References

23. Kral, R. 1992. A new species of *Fimbristylis* (Cyperaceae) from the sandstone and granitic outcrops of Alabama and Georgia. *Sida* 15: 317-321.
24. Kükenthal, G. 1935-1936. Cyperaceae-Scirpoideae-Cypereae. In A. Engler, ed., *Pflanzenreich* IV. 20 (Heft) 101:1-671.
25. Tucker, G. C. 1987. The genera of Cyperaceae in the southeastern United States. *J. Arnold Arbor.* 68:361-445.
26. Bruhl, J. 1993. *Sedge Genera of the World (Version 2.01)*, Intkey, Interactive Dataset. N.C.W. Beadle Herbarium, Univ. of New England, Armidale. New South Wales, Australia.
27. Soros, C. L. and Bruhl, J. J. 2000. Multiple Evolutionary Origins of C4 Photosynthesis in the Cyperaceae. Pages 629-636, in K. L. Wilson and D. A. Morrison, *Monocots: Systematics and Evolution*. CSIRO Publishing. Collingwood, Victoria, Australia.
28. Bryson, C. T., R. Carter, L. B. McCarty, and F. H. Yelverton. 1997. *Kyllinga*, a genus of neglected weeds in the continental United States. *Weed Technol.* 11: 838-842.
29. Michaux, A. 1803. *Flora boreali-americana*. Vol. I. Paris.
30. Elliott, S. 1821. *A sketch of the botany of South-Carolina and Georgia*. Vol. 1. J. R. Schenck. Charleston, South Carolina.
31. Torrey, J. 1836. Monograph of North American Cyperaceae. *Ann. Lyceum Nat. Hist. New York* 3: 249-288.
32. Wagner, W. L., D. R. Herbst and S. H. Sohmer. 1990. *Manual of the flowering plants of Hawaii*. Vol. 2. University of Hawaii Press, Bishop Museum. Honolulu.
33. Wagner, W. L. and D. R. Herbst. 1995. Contributions to the flora of Hawaii. IV. New records and name changes. *Bishop Mus. Occas. Pap.* 42: 13-27.